



TITLE:

Mapping theorems for SC -spaces (General and geometric topology today and their problems)

AUTHOR(S):

Komoda, Chieko

CITATION:

Komoda, Chieko. Mapping theorems for SC -spaces (General and geometric topology today and their problems). 数理解析研究所講究録 2008, 1578: 19-21

ISSUE DATE:

2008-02

URL:

<http://hdl.handle.net/2433/81380>

RIGHT:

Mapping theorems for C -spaces

順天堂大学スポーツ健康科学部

菰田智恵子 (Chieko Komoda)

We assume that all spaces are normal unless otherwise stated. We refer the readers to [2] for dimension theory.

In this note we study mapping theorems for C -spaces.

A space X is a C -space (an A -weakly infinite-dimensional space) if for every countable collection $\{\mathcal{G}_i : i \in \mathbb{N}\}$ of open covers (two-element open covers, respectively) of X there exists a countable collection $\{\mathcal{H}_i : i \in \mathbb{N}\}$ of collections of pairwise disjoint open subsets of X such that $\mathcal{H}_i < \mathcal{G}_i$ for every $i \in \mathbb{N}$ and $\bigcup_{i=1}^{\infty} \mathcal{H}_i$ covers X (cf. [1]).

Evidently, every C -space is A -weakly infinite-dimensional. However, it is not known whether the converse is true.

Polkowski [5] proved the following theorem.

Theorem 1 (Polkowski [5]). *If $f : X \longrightarrow Y$ is a closed mapping of an A -weakly infinite-dimensional countably paracompact space X onto a space Y and there exists an integer $k \geq 1$ such that $|f^{-1}(y)| \leq k$ for every $y \in Y$, then Y is A -weakly infinite-dimensional.*

We proved that the following theorem, which is an analogous result for C -spaces.

Theorem 2. *If $f : X \longrightarrow Y$ is a closed mapping of a countably paracompact C -space X onto a paracompact space Y and there exists an integer $k \geq 1$ such that $|f^{-1}(y)| \leq k$ for every $y \in Y$, then Y is a C -space.*

Problem. Does theorem 1 (or theorem 2) hold for closed mappings with finite fibers?

In [4], Pol proved the following theorem.

Theorem 3 (Pol [4]). *If $f : X \longrightarrow Y$ is a continuous mapping of a compact metrizable space X onto a metrizable space Y such that $|f^{-1}(y)| \leq \aleph_0$ for every $y \in Y$, then X is an A -weakly infinite-dimensional space (resp. a C -space) if and only if Y is an A -weakly infinite-dimensional space (resp. a C -space).*

Does Theorem 3 remain true if we replace ' $|f^{-1}(y)| \leq \aleph_0$ ' by ' $|f^{-1}(y)| < c$ '? In [5], Polkowski proved the following theorem.

Theorem 4 (Polkowski [5]). *If $f : X \longrightarrow Y$ is a continuous mapping of a compact A -weakly infinite-dimensional space X onto a space Y such that $|f^{-1}(y)| < c$ for every $y \in Y$, then Y is A -weakly infinite-dimensional.*

Similarly, the following theorem holds.

Theorem 5. *If $f : X \longrightarrow Y$ is a continuous mapping of a compact C -space X onto a space Y such that $|f^{-1}(y)| < c$ for every $y \in Y$, then Y is a C -space.*

On the other hand, Hattori and Yamada proved that the following theorem.

Theorem 6 (Hattori and Yamada [3]).

- (i) *If $f : X \longrightarrow Y$ is a closed mapping of a countably paracompact (or hereditarily normal) space X onto a C -space Y such that $f^{-1}(y)$ is A -weakly infinite-dimensional for every $y \in Y$, then X is A -weakly infinite-dimensional.*
- (ii) *If $f : X \longrightarrow Y$ is a closed mapping of a paracompact space X onto a C -space Y such that $f^{-1}(y)$ is a C -space for every $y \in Y$, then X is a C -space.*

Problem. Does Theorem 6(i) remain true if we replace ' $f^{-1}(y)$ is a C -space' by ' $f^{-1}(y)$ is A -weakly infinite-dimensional'?

References

- [1] D. F. Addis and J. H. Gresham, *A class of infinite-dimensional spaces, Part I: Dimension theory and Alexandroff's Problem*, Fund. Math. 101(1978), 195-205.
- [2] R. Engelking, *Theory of Dimensions, Finite and Infinite*, Heldermann Verlag, 1995.
- [3] Y. Hattori and K. Yamada, *Closed pre-images of C-spaces*, Math. Japonica 34(1989), 555-561.
- [4] R. Pol, *On light mappings without perfect fibers on compacta*, Tsukba J. Math. 20(1996), 11-19.
- [5] L. Polkowski, *Some theorems on invariance of infinite dimension under open and closed mappings*, Fund. Math. 119(1983), 11-34.

Chieko Komoda

Department of Health Science, School of Health & Sports Science, Juntendo University
Inba, Chiba, 270-1695, Japan

E-mail address: chieko_komoda@sakura.juntendo.ac.jp